FREUD'S EQUATIONS FOR ORTHOGONAL POLYNOMIALS AS DISCRETE PAINLEVÉ EQUATIONS.

Alphonse P. Magnus

ABSTRACT. We consider orthogonal polynomials p_n with respect to an exponential weight function $w(x) = \exp(-P(x))$. The related equations for the recurrence coefficients have been explored by many people, starting essentially with Laguerre [49], in order to study special continued fractions, recurrence relations, and various asymptotic expansions (G. Freud's contribution [28,56]).

Most striking example is $n = 2tw_n + w_n(w_{n+1} + w_n + w_{n-1})$ for the recurrence coefficients $p_{n+1} = xp_n - w_n p_{n-1}$ of the orthogonal polynomials related to the weight $w(x) = \exp(-4(tx^2 + x^4))$ (notations of [26, pp.34–36]). This example appears in practically all the references below. The connection with discrete Painlevé equations is described here.

1. CONSTRUCTION OF ORTHOGONAL POLYNOMIALS RECURRENCE COEFFICIENTS.

Consider the set $\{p_n\}_0^\infty$ of orthonormal polynomials with respect to a weight function w on (a part of) the real line:

$$\int_{-\infty}^{\infty} p_n(x)p_m(x)w(x)\,dx = \delta_{n,m}, \qquad n, m = 0, 1, \ldots \tag{1}$$

The very useful *recurrence formula* has the form $a_1 p_1(x) = (x - b_0)p_0$,

$$a_{n+1}p_{n+1}(x) = (x - b_n)p_n(x) - a_n p_{n-1}(x), \quad n = 1, 2, \ldots \tag{2}$$

The connection between the nonnegative integrable function w and the real sequences $\{a_n\}_1^\infty, \{b_n\}_0^\infty$ is of the widest interest. It is investigated in combinatorics [9,10,52], asymptotic analysis [11,12,28,30,53–55,61,77–79], numerical analysis [15, 21, 31, 32, 51, 57], and, of course, Lax-Painlevé-Toda theory (all the other references, excepting *perhaps* Chaucer).

However, the connection may seem quite elementary and explicit:

1. From w to a_n and b_n: let $\mu_k = \int_{-\infty}^{\infty} x^k\, w(x)\,dx$, $k = 0, 1, \ldots$ be the moments of w, then

$$a_n^2 = \frac{H_{n-1}H_{n+1}}{(H_n)^2}, n = 1, 2, \ldots \qquad b_n = \frac{\widetilde{H}_{n+1}}{H_{n+1}} - \frac{\widetilde{H}_n}{H_n}, n = 0, 1, \ldots \tag{3}$$

where H_n and $\widetilde{H}_n$ are the determinants

$$H_n = \begin{vmatrix} \mu_0 & \cdots & \mu_{n-1} \\ \vdots & & \vdots \\ \mu_{n-1} & \cdots & \mu_{2n-2} \end{vmatrix} \quad ; \widetilde{H}_n = \begin{vmatrix} \mu_0 & \cdots & \mu_{n-2} & \mu_n \\ \vdots & & \vdots & \vdots \\ \mu_{n-1} & \cdots & \mu_{2n-3} & \mu_{2n-1} \end{vmatrix}, \qquad (4)$$

$n = 0, 1, \ldots, H_0 = 1, \widetilde{H}_0 = 0$.

2. From the a_n's and b_n's to w: consider the Jacobi matrix

$$\boldsymbol{J} = \begin{bmatrix} b_0 & a_1 & & & \\ a_1 & b_1 & a_2 & & \\ & a_2 & b_2 & a_3 & \\ & & \ddots & \ddots & \ddots \end{bmatrix}$$

then, remarking that (2) may be written $xp_n(x) = a_n p_{n-1}(x) + b_n p_n(x) + a_{n+1} p_{n+1}(x)$, (Remark that the coefficients appear now with positive signs, a very elementary fact, but always able to puzzle weak souls), or $x\,\boldsymbol{p}(x) = \boldsymbol{J}\boldsymbol{p}(x)$, where $\boldsymbol{p}(x)$ is the column (infinite) vector of $p_0(x), p_1(x), \ldots$, so that the expansion of $x^k p_n(x)$ in the basis $\{p_0, p_1, \ldots\}$ is $x^k p_n(x) = \sum_\ell (\boldsymbol{J}^k)_{n,\ell} p_\ell(x)$. As we are dealing with a basis of orthonormal polynomials, this means that $(\boldsymbol{J}^k)_{n,\ell}$ is the ℓ^{th} Fourier coefficient of $x^k p_n(x)$, (where row and column indexes start at zero), i.e., $(\boldsymbol{J}^k)_{n,\ell} = \int_{-\infty}^{\infty} x^k p_n(x) p_\ell(x) w(x)\,dx$. For any polynomial p, one has

$$(p(\boldsymbol{J}))_{n,\ell} = \int_{-\infty}^{\infty} p(x) p_n(x) p_\ell(x) w(x)\,dx. \qquad (5)$$

In particular, $(\boldsymbol{J}^k)_{0,0} = \mu_k/\mu_0$.

Direct numerical use of (3) is almost always unsatisfactory, for stability reasons. Together with various ways to cope with the numerical stability problem [31,32], compact formulas or equations for the recurrence coefficients a_n and b_n in special cases have been sought. Of course, quite a number of exact solutions are known (appendix of [14]) and new ones are steadily discovered (Askey-Wilson polynomials and other instances of the new "$q-$calculus" [4, 35,36,47]). The trend explored here is essentially related to weight functions satisfying simple differential equations. Main innovators were Laguerre in 1885 [49], Shohat in 1939 [77], and Freud in 1976 [28]. Their contributions are now described in reverse order. On Géza Freud (1922-1979), see [29,68,69].

2. Definition of Freud's equations.

In order to reduce the technical contents of what will follow, only *even* weights: $\forall$ real x, $\quad w(-x) = w(x)$ will be used. Then, the orthogonal polynomials are even or odd according to their degrees; equivalently, all the b_n's in (2), in (3), and in the Jacobi matrix $\boldsymbol{J}$ vanish. Let $\boldsymbol{a}$ be the sequence of coefficients $\{a_1, a_2, \ldots\}$.

Theorem 1. *Let P be the real even polynomial $P(x) = c_0 x^{2m} + c_1 x^{2m-2} + \cdots + c_m$, with $c_0 > 0$, and let $a_1, a_2, \ldots$ be the recurrence coefficients* (2) *of*

the orthonormal polynomials with respect to the weight $w(x) = \exp(-P(x))$ *on the whole real line. Then, the* **Freud's equations**

$$F_n(\boldsymbol{a}) := a_n \left(P'(\boldsymbol{J})\right)_{n,n-1} = n, \quad n = 1, 2, \ldots \tag{6}$$

hold, where $(\ )_{n,n-1}$ means the element at row n and column $n-1$ of the (infinite) matrix $P'(\boldsymbol{J})$. A finite number of computations is involved in each of these elements. The row and column indexes start at 0. We could as well take the indexes $(n-1, n)$, as $\boldsymbol{J}$ and any polynomial function of $\boldsymbol{J}$ are symmetric matrices.

Remark. One has

$$F_1(\boldsymbol{a}) = \left(\boldsymbol{J}P'(\boldsymbol{J})\right)_{0,0}. \tag{7}$$

Indeed, the first element of the first row of $\boldsymbol{J}P'(\boldsymbol{J})$ is a_1 times the element $(1,0)$ of $P'(\boldsymbol{J})$, i.e., $F_1(\boldsymbol{a})$.

Examples.

$$w(x) = \exp(-x^2) \quad F_n(\boldsymbol{a}) = 2a_n^2, \tag{8}$$

$$w(x) = \exp(-\alpha x^4 - \beta x^2) \quad F_n(\boldsymbol{a}) = 4\alpha a_n^2(a_{n-1}^2 + a_n^2 + a_{n+1}^2) + 2\beta a_n^2, \tag{9}$$

$$\begin{aligned} w(x) = \exp(-x^6) \quad F_n(\boldsymbol{a}) &= 6a_n^2(a_{n-2}^2 a_{n-1}^2 + a_{n-1}^4 + 2a_{n-1}^2 a_n^2 + a_n^4 \\ &\quad + a_{n-1}^2 a_{n+1}^2 + 2a_n^2 a_{n+1}^2 + a_{n+1}^4 + a_{n+1}^2 a_{n+2}^2), \end{aligned} \tag{10}$$

The elementary Hermite polynomials case is of course immediately recovered in (8). (9) and (10) were used by Freud [28] in investigations on asymptotic behaviour, see [11,53,56,57,67,69] for more. The rich connection of (9) with discrete and continuous Painlevé theory will be recalled later on. For a much more general setting, see [1, Theor. 4.1].

Proof of (6). We consider two different ways to write the integral $I_n = \displaystyle\int_{-\infty}^{\infty} p_n'(x) p_{n-1}(x) \exp(-P(x))\, dx$. First, let $p_{n-1}(x) = \pi_{n-1}x^{n-1} + \ldots$ and $p_n(x) = \pi_n x^n + \ldots$. From (2) (with $n-1$ instead of n), $\pi_n = \pi_{n-1}/a_n$; so, $p_n'(x) = n\pi_n x^{n-1} + \ldots = \dfrac{n}{a_n} p_{n-1}(x) +$ a polynomial of degree $\leqslant n-2$. As p_{n-1} is orthogonal to all polynomials of degree $\leqslant n-2$, what remains is $I_n = \dfrac{n}{a_n}$, as the orthonormal polynomials have a unit square integral. Next, we perform an integration by parts on $I_n = -\displaystyle\int_{-\infty}^{\infty} p_n(x) \left[p_{n-1}(x) \exp(-P(x))\right]' dx = \displaystyle\int_{-\infty}^{\infty} P'(x) p_n(x) p_{n-1}(x) \exp(-P(x))\, dx$, using the orthogonality of p_n and p_{n-1}' of degree $< n-1 < n$, and the latter integral is $(P'(\boldsymbol{J}))_{n,n-1}$, according to the spectral representation (5). □

3. Freud's equations software.

The interested reader (if there is still one left) will find in the "software" part of http://www.math.ucl.ac.be/~magnus/ three FORTRAN programs related to Freud's equations. `freud1.f` asks for an exponent m and puts data associated to the Freud's equations for $x^2, x^4, \ldots, x^{2m}$ in the file `freud00`m`.dat`. `freud2.f` produces a readable (sort of) listing out of such a data file. Finally, `freud3.f` reads the coefficients of a polynomial P and computes in a stable way a sequence of positive recurrence coefficients $a_1, a_2, \ldots$ associated to the weight $\exp(-P(x))$ on the whole real line. Actually, weight functions $|x|^\alpha \exp(-P(x))$, with P even, on the whole real line, and $x^\beta \exp(-Q(x))$ on the *positive* real line are considered simultaneously ($P(x) = Q(x^2)$).

4. Solution of Freud's equations.

We started from the *solution* (3), and built the *equation* (6) afterwards, when the a_n's are recurrence coefficients of orthogonal polynomials associated to the weight $\exp(-P(x))$.

We try now to discuss the general solution of (6). With P even of degree $2m$, the n^{th} equation of (6) involves $a_{n-m+1}, \ldots, a_{n+m-1}$ (see [56, 57] or the examples (8) (9), (10)), so that the general solution depends on $2m-2$ arbitrary constants, for instance $a_{n_0-m+1}, \ldots, a_{n_0+m-2}$. I shall only solve the case $a_{-m+1} = \cdots = a_0 = 0$, and investigate how the solution depends on the $m-1$ initial data $a_1, \ldots, a_{m-1}$:

Theorem 2. *With $P(x) = c_0 x^{2m} + c_1 x^{2m-2} + \cdots + c_m$, the solution of (6) with $a_{-m+1} = \cdots = a_0 = 0$ is (3), with (4), where the μ_j's of (4) satisfy the* linear *recurrence relation of order $2m$*

$$2mc_0\mu_{2n+2m} + (2m-2)c_1\mu_{2n+2m-2} + \cdots + 2c_{m-1}\mu_{2n+2} = (2n+1)\mu_{2n}, \quad (11)$$

for $n = 0, 1, \ldots$, and $\mu_{2j+1} = 0$. $\mu_2, \mu_4, \ldots, \mu_{2m-2}$ are given by $\mu_{2j}/\mu_0 = (\boldsymbol{J}^{2j})_{0,0}$.

The following lemma will be used:

Lemma 1. *Let $\boldsymbol{a}^{(k)} = \{a_{n,k}\}_{n=1}^{\infty}$, $k = 0, 1, \ldots$, be the solutions of the* ***quotient-difference*** *equations*

$$\begin{aligned} a_{n-1,k+1}^2 + a_{n,k+1}^2 &= a_{n,k}^2 + a_{n+1,k}^2, && \text{if } n \text{ is odd,} \\ a_{n-1,k+1}a_{n,k+1} &= a_{n,k}a_{n+1,k}, && \text{if } n \text{ is even,} \end{aligned} \quad (12)$$

then, $F_n(\boldsymbol{a}^{(k)})$ satisfies

$$\begin{aligned} \frac{a_{n-1,k+1}}{a_{n,k}}\left(F_n(\boldsymbol{a}^{(k+1)}) - F_n(\boldsymbol{a}^{(k)})\right) &= \frac{a_{n,k}}{a_{n-1,k+1}}\left(F_{n+1}(\boldsymbol{a}^{(k)}) - F_{n-1}(\boldsymbol{a}^{(k+1)})\right), \\ & && n \text{ even,} \\ F_n(\boldsymbol{a}^{(k+1)}) - F_n(\boldsymbol{a}^{(k)}) &= F_{n+1}(\boldsymbol{a}^{(k)}) - F_{n-1}(\boldsymbol{a}^{(k+1)}), && n \text{ odd.} \end{aligned} \quad (13)$$

where $F_n(\boldsymbol{a}^{(k)}) = a_{n,k}\,(P'(\boldsymbol{J}_k))_{n,n-1}$, *and where* $\boldsymbol{J}_k$ *is the Jacobi matrix*

$$\boldsymbol{J}_k = \begin{bmatrix} 0 & a_{1,k} & & & \\ a_{1,k} & 0 & a_{2,k} & & \\ & a_{2,k} & 0 & a_{3,k} & \\ & & \ddots & \ddots & \ddots \end{bmatrix}$$

The quotient-difference equations are well known in orthogonal polynomials identities investigations, see [74, 75] for recent examples. The notation chosen here is motivated by the interpretation of $a_{n,k}$ as recurrence coefficient of orthogonal polynomials related to the weight $x^{2k}w(x)$, but we do not need this interpretation in order to establish the theorem and the lemma. Actually, the theorem 2 is meant to recover an interpretation of the solutions of (6) in terms of orthogonal polynomials, but arbitrary initial data $a_1, a_2, \ldots, a_{m-1}$ will yield formal orthogonal polynomials through formal moments μ_j which, from (11), can still be written $\mu_j = \int_S x^j \exp(-P(x))\,dx$, but where the support S is now a system of arcs in the complex plane, see [66, "Laplace's method"], [62,63].

Remark that, with $a_{n,0} = a_n$ and $a_{0,k} = 0$, all the $a_{n,k}$'s are completely determined by (12) as functions of $a_1, a_2, \ldots$

Proof of Lemma 1. The quotient-difference equations (12) can be written $\boldsymbol{J}_k\boldsymbol{L}_k = \boldsymbol{L}_k\boldsymbol{J}_{k+1}$, where $\boldsymbol{L}_k$ is the infinite lower triangular matrix

$$\boldsymbol{L}_k = \begin{bmatrix} a_{1,k} & & & & & \\ 0 & a_{1,k+1} & & & & \\ a_{2,k} & 0 & a_{3,k} & & & \\ & a_{2,k+1} & 0 & a_{3,k+1} & & \\ & & a_{4,k} & 0 & a_{5,k} & \\ & & & \ddots & \ddots & \ddots \end{bmatrix}$$

One finds also $\boldsymbol{J}_k^2 = \boldsymbol{L}_k\boldsymbol{R}_k$, $\boldsymbol{J}_{k+1}^2 = \boldsymbol{R}_k\boldsymbol{L}_k$, where $\boldsymbol{R}_k$ is the transposed of $\boldsymbol{L}_k$ (the famous LR relations). So, $\boldsymbol{J}_k^3\boldsymbol{L}_k = \boldsymbol{L}_k\boldsymbol{R}_k\boldsymbol{L}_k\boldsymbol{J}_{k+1} = \boldsymbol{L}_k\boldsymbol{J}_{k+1}^3$, $\ldots$, $\boldsymbol{J}_k^{2p+1}\boldsymbol{L}_k = \boldsymbol{L}_k\boldsymbol{R}_k\boldsymbol{L}_k\boldsymbol{J}_{k+1}^{2p-1} = \boldsymbol{L}_k\boldsymbol{J}_{k+1}^{2p+1}$, for any odd power, so $P'(\boldsymbol{J}_k)\boldsymbol{L}_k = \boldsymbol{L}_kP'(\boldsymbol{J}_{k+1})$, and we look at the element of indexes $(n, n-1)$:

$$(P'(\boldsymbol{J}_k))_{n,n-1}(\boldsymbol{L}_k)_{n-1,n-1} + (P'(\boldsymbol{J}_k))_{n,n+1}(\boldsymbol{L}_k)_{n+1,n-1} = (\boldsymbol{L}_k)_{n,n-2}(P'(\boldsymbol{J}_{k+1}))_{n-2,n-1} + (\boldsymbol{L}_k)_{n,n}(P'(\boldsymbol{J}_{k+1}))_{n,n-1},$$

$$\text{or}\quad F_n(\boldsymbol{a}^k)\frac{(\boldsymbol{L}_k)_{n-1,n-1}}{a_{n,k}} + F_{n+1}(\boldsymbol{a}^k)\frac{(\boldsymbol{L}_k)_{n+1,n-1}}{a_{n+1,k}} = \frac{(\boldsymbol{L}_k)_{n,n-2}}{a_{n-1,k+1}}F_{n-1}(\boldsymbol{a}^{k+1}) + \frac{(\boldsymbol{L}_k)_{n,n}}{a_{n,k+1}}F_n(\boldsymbol{a}^{k+1}),$$

leading to (13). □

Proof of Theorem 2. From (13), with (6), ($\boldsymbol{a}^{(0)} = \boldsymbol{a}$), and $F_0(\boldsymbol{a}^{(k)}) = 0$ for all k, one finds by induction $F_n(\boldsymbol{a}^{(k)}) = n + k(1-(-1)^n)$, so, using (7), $F_1(\boldsymbol{a}^{(k)}) = (\boldsymbol{J}_kP'(\boldsymbol{J}_k))_{0,0} = 2k+1$, $k = 0, 1, \ldots$ Let μ_{2k} = constant $\times a_1^2a_{1,1}^2a_{1,2}^2\cdots a_{1,k-1}^2$ and $\mu_{2k+1} = 0$. From the LR relations, $(\boldsymbol{J}_k^{2p})_{0,0} = (\boldsymbol{L}_k\boldsymbol{L}_{k+1}\cdots\boldsymbol{L}_{k+p}\boldsymbol{R}_{k+p}\cdots\boldsymbol{R}_{k+1}\boldsymbol{R}_k)_{0,0} = a_{1,k}^2a_{1,k+1}^2\cdots a_{1,k+p-1}^2 = \mu_{2k+2p}/\mu_{2k}$.

Finally, $2k+1 = (\boldsymbol{J}_k P'(\boldsymbol{J}_k))_{0,0} = 2mc_0(\boldsymbol{J}_k^{2m})_{0,0} + (2m-2)c_1(\boldsymbol{J}_k^{2m-2})_{0,0} + \cdots + 2c_{m-1}(\boldsymbol{J}_k^2)_{0,0} = 2mc_0\mu_{2k+2m}/\mu_{2k} + (2m-2)c_1\mu_{2k+2m-2}/\mu_{2k} + \cdots + 2c_{m-1}\mu_{2k+2}/\mu_{2k}$, whence (11), and the determinant ratios (3) follow from (12): $a_{n,k}^2 = H_{n-1,k}H_{n+1,k}/(H_{n,k})^2$ where $H_{n,k}$ is the Hankel determinant of rows $\mu_{2k}, \mu_{2k+1}(=0), \ldots, \mu_{2k+n-1}$; $\mu_{2k+1}, \ldots, \mu_{2k+n}$, etc., up to $\mu_{2k+n-1}, \ldots, \mu_{2k+2n-2}$. □

The *isomonodromy approach* looks at the linear differential equation satisfied by an orthogonal polynomial p_n [5, 16, 17, 37, 38] and produces isomonodromy identities of interest here. For instance, working (9) leads to $a_n^2 = (4\alpha)^{-1/2}\mathrm{P}_{\mathrm{IV}}((4\alpha)^{-1/2}\beta; -n/2, -n^2/4)$ (Kitaev, [23–26,43]), where $\mathrm{P}_{\mathrm{IV}}(t; A, B)$ is the solution of the fourth Painlevé equation

$$\ddot{y} = \frac{\dot{y}^2}{2y} + \frac{3y^3}{2} + 4ty^2 + 2(t^2 - A)y + \frac{B}{y}$$

which remains $O(t^{-1})$ when $t \to +\infty$ [60, pp. 231-232] (the importance and relevance of this latter condition is not yet quite clear). See [1–3, 16–20, 22–26, 39, 43, 44, 46] for more on such connections.

5. Freud's equations as discrete Painlevé equations.

As far as I understand the matter, discrete Painlevé equations were first designed as clever discretisations of the genuine Painlevé equations, so to keep interesting features of these equations, especially integrability. For instance, (9) when $\beta \neq 0$ is a discretisation of the Painlevé-I equation [23–26, 33, 34, 43]. Then, these features are examined on various discrete equations not necessarily associated to Painlevé equations. So, for instance, [33, p.350] encounter (9) with $\beta = 0$ as a discrete equation which is no more linked to the discretisation of a continuous Painlevé equation (and the authors of [33] call (9) a discrete Painlevé-0 equation when $\beta = 0$).

If we want to check that the Freud's equations are a valuable instance of discrete Painlevé equation, the following features are of interest, according to experts [48]:

5.1. **Analyticity.** Each a_{n+m-1}^2 is a rational function of preceding elements, so a meromorphic function of initial conditions.

5.2. **Reversibility.** There is no way to distinguish past and future, (6) is left unchanged when $a_{n+1}, a_{n+2}, \ldots$ are permuted with $a_{n-1}, a_{n-2}, \ldots$ Indeed ([56, p.369]), (6) is a sum of terms $a_n^2 a_{n+i_1}^2 a_{n+i_2}^2 \cdots a_{n+i_p}^2$, provided $0 \leqslant i_1+1 \leqslant i_2+2 \leqslant \cdots \leqslant i_p+p \leqslant p+1$. Then, $\{-i_1, -i_2, \ldots, -i_p\}$ satisfies the same conditions and is therefore present too: $0 \leqslant -i_p+1 \leqslant -i_{p-1}+2 \leqslant \cdots \leqslant -i_1+p \leqslant p+1$.

5.3. **Symmetry.** An even stronger property is the following:

Theorem 3. *The matrix of derivatives* $\left(\dfrac{\partial F_n(\boldsymbol{a})}{\partial \log a_m}\right)_{n,m=1,2,\ldots}$ *is symmetric.*

Proof: see [56,57].

5.4. **Integrability.** The existence of the formula (3) according to Theorem 2 shows how (6) can be solved. The similarity of (3) with known solutions of discrete Painlevé equations is also striking [42, 45]. The structure of the solution leads also to

5.5. **Singularity confinement.** The movable poles property of continuous equations is replaced by the following [34]: if initial values are such that some a_n, say a_{n_0} becomes very large, only a finite number of neighbours $a_{n_0+1}, \ldots, a_{n_0+p}$ should be liable to be very large too; moreover, a_{n_0+p+1}, etc. should be continuous functions of a_{n_0-1}, a_{n_0-2} etc. The formula (3) shows that the singularities of a_n are the zeros of H_n which are determinants built with solutions of (11), therefore continuous functions of the initial data.

6. Generalizations of Freud's equations: semiclassical orthogonal polynomials, etc.

Let the weight function satisfy the differential equation $w'/w = 2V/W$, where V and W are polynomials. The related orthogonal polynomials on a support S are then called *semi-classical* [6–8, 40, 41, 64, 65, 72] (classical: degrees of V and $W \leqslant 1$ and 2). One can still build equations for the recurrence coefficients by working out the integrals $I_n = \displaystyle\int_S W(x)p_n'(x)p_{n-1}(x)w(x)\,dx$, and $J_n = \displaystyle\int_S W(x)p_n'(x)\; p_n(x)w(x)\,dx$. Many equivalent forms can be found, and it not yet clear to know which one is the most convenient. If we manage to have $W(x)w(x) = 0$ at the endpoints of the support S, explicit formulas follow [77] (see also formula (4.5) of [1]). For instance, with $w(x) = \exp(-Ax^2)$ on $S = [-a, a]$ (see [15] for other equations), one takes $W(x) = x^2 - a^2$ and $V(x) = -Ax(x^2 - a^2)$, uses $p_n' = np_{n-1}/a_n + [2(a_1^2 + \cdots + a_{n-1}^2) - na_{n-1}^2]p_{n-3}/(a_{n-2}a_{n-1}a_n) + \cdots$ from (2), to find $2(a_1^2 + \cdots + a_{n-1}^2) + (2n + 1)a_n^2 - na^2 - 2Aa_n^2(a_{n-1}^2 + a_n^2 + a_{n+1}^2 - a) = 0$, or $(2n + 1)x_{n+1} - (2n - 1)x_n - na^2 - 2A(x_{n+1} - x_n)(x_{n+2} - x_{n-1} - a) = 0$, where $x_n = a_1^2 + \cdots + a_{n-1}^2$.

It is not clear how to recover reversibility and symmetry. For an even much more nasty case, see [61].

But the root of the matter lies even deeper, as already recognized by Laguerre in 1885 [49]! The point is to derive equations for the coefficients in the Jacobi continued fraction $f(z) = \cfrac{1}{z - b_0 - \cfrac{a_1^2}{z - b_1 - \cdots}}$ from a differential equation $Wf' = 2Vf + U$ with polynomial coefficients (same W and V as above: orthogonal polynomials lead to this continued fraction through $f(z) = \displaystyle\int_S (z - x)^{-1}w(x)\,dx$). See [30, 72] for this technique.

The continued fraction approach leads readily to the *Riccati* extension $Wf' = Rf^2 + 2Vf + U$ [20,55], as each continued fraction $f_k(z) = \cfrac{1}{z - b_k - \cfrac{a_{k+1}^2}{z - b_{k+1} - \cdots}}$ is found to satisfy a Riccati equation $Wf_k' = -a_k^2\Theta_{k-1}f_k^2 + 2\Omega_k f_k - \Theta_k$ too, where Θ_k and Ω_k are polynomials of fixed degrees (i.e., independent. of k). Equations for the recurrence coefficients a_k's and b_k's follow from recurrence relations for the Θ_k's and Ω_k's (use $f_0(z) = f(z)$ and $f_k(z) = 1/[z - b_k - a_{k+1}^2 f_{k+1}(z)]$).

Finally, the differential operator can be replaced by a suitable difference operator, amounting to $W(z)\dfrac{f(\varphi_2(z)) - f(\varphi_1(z))}{\varphi_2(z) - \varphi_1(z)} = R(z)f(\varphi_1(z))f(\varphi_2(z)) + V(z)[f(\varphi_1(z)) + f(\varphi_2(z))] + U(z)$, where $\varphi_1(z)$ and $\varphi_2(z)$ are the two roots of a quadratic equation $A\varphi^2(z) + 2Bz\varphi(z) + Cz^2 + 2D\varphi(z) + 2Ez + F = 0$, see [36,58].

Acknowledgements.

Thanks to the Conference organizers, and to Darwin's College hospitality, according to the highest quality standards since more than 600 years:

26: In felaweshipe, and pilgrimes were they alle,	In fellowship, and pilgrims were they all
27: That toward caunterbury wolden ryde.	That toward Canterbury town would ride.
28: The chambres and the stables weren wyde,	The rooms and stables spacious were and wide,
29: And wel we weren esed atte beste.	And well we there were eased, and of the best.

University of Virginia Library Electronic Text Center. : http://etext.virginia.edu/mideng.browse.html, Cha2Can
Available from: Oxford Text Archive no. OTA U-1678-C 1993 [13]

Linkname: Canterbury Tales (prologue)
URL: gopher://gopher.vt.edu:10010/02/63/38

References

[1] M. Adler, P. van Moerbeke, Matrix integrals, Toda symmetries, Virasoro constraints, and orthogonal polynomials, *Duke Math. J.* **80** (1995) 863-911.

[2] A.I. APTEKAREV, Asymptotic properties of polynomials orthogonal on a system of contours and periodic motions of Toda lattices, *Mat. Sb.* **125** (1984) 231-258, = *Math. USSR Sbornik* **53** (1986) 233-260.

[3] A.I. Aptekarev, A. Branquinho, F. Marcellán, Toda-type differential equations for the recurrence coefficients of orthogonal polynomials and Freud transformation, Pré-Publicações Dep. Matemática Univ. Coimbra, February 1996.
[4] N.M. Atakishiyev, These Proceedings.
[5] F.V.ATKINSON, W.N.EVERITT, Orthogonal polynomials which satisfy second order differential equations, pp. 173-181 *in E.B.Christoffel* (P.L.BUTZER and F.FEHÉR, editors), Birkhäuser, Basel, 1981.
[6] S. BELMEHDI, On semi-classical linear functionals of class $s = 1$. Classification and integral representations. *Indag. Mathem.* N.S. **3** (3) (1992), 253-275.
[7] S. BELMEHDI, A. RONVEAUX, Laguerre-Freud's equations for the recurrence coefficients of semi-classical orthogonal polynomials, *J. Approx. Theory* **76** (1994) 351-368.
[8] S. BELMEHDI, A. RONVEAUX, On the coefficients of the three-term recurrence relation satisfied by some orthogonal polynomials. *Innovative Methods in Numerical Analysis*, Bressanone, Sept. 7-11$^{\text{th}}$, 1992.
[9] D. BESSIS, A new method in the combinatorics of the topological expansion, *Comm. Math. Phys.* **69** (1979), 147-163.
[10] D. BESSIS, C. ITZYKSON, J.B. ZUBER, Quantum field theory techniques in graphical enumeration, *Adv. in Appl. Math.* **1** (1980), 109-157.
[11] S. BONAN, P.NEVAI, Orthogonal polynomials and their derivatives,I, *J. Approx. Theory* **40** (1984), 134-147.
[12] S.S. BONAN, D.S. LUBINSKY, P.NEVAI, Orthogonal polynomials and their derivatives,II, *SIAM J. Math. An.* **18** (1987), 1163-1176.
[13] G. CHAUCER, *The works of Geoffrey Chaucer* 2nd. ed. editor: F.N. Robinson. : Houghton Mifflin. Boston, Mass. 1957
[14] T. Chihara, *An Introduction to Orthogonal Polynomials*, Gordon & Breach, 1978.
[15] R.C.Y. CHIN, A domain decomposition method for generating orthogonal polynomials for a Gaussian weight on a finite interval, *J. Comp. Phys.* **99** (1992) 321-336.
[16] D.V. CHUDNOVSKY, Riemann monodromy problem, isomonodromy deformation eqations and completely integrable systems, pp.385-447 *in Bifurcation Phenomena in Mathematical Physics and Related Topics, Proceedings Cargèse, 1979* (C.BARDOS & D.BESSIS, editors), NATO ASI series C, vol. **54**, D.Reidel, Dordrecht, 1980.
[17] G.V. CHUDNOVSKY, Padé approximation and the Riemann monodromy problem, pp.449-510 *in Bifurcation Phenomena in Mathematical Physics and Related Topics, Proceedings Cargèse, 1979* (C.BARDOS & D.BESSIS, editors), NATO ASI series C, vol. **54**, D.Reidel, Dordrecht, 1980.
[18] D.V.CHUDNOVSKY, G.V.CHUDNOVSKY, Laws of composition of Bäcklund transformations and the universal form of completely integrable systems in dimensions two and three, *Proc. Nat. Acad. Sci. USA* **80** (1983) 1774-1777.
[19] D.V.CHUDNOVSKY, G.V.CHUDNOVSKY, High precision computation of special function in different domains, talk given at Symbolic Mathematical Computation conference, Oberlech, July 1991.
[20] D.V. Chudnovsky, G.V. Chudnovsky, Explicit continued fractions and quantum gravity, *Acta Applicandæ Math.* **36** (1994) 167–185.
[21] A.S. CLARKE, B. SHIZGAL, On the generation of orthogonal polynomials using asymptotic methods for recurrence coefficients. *J. Comp. Phys.* **104** (1993) 140-149.
[22] P. Deift, K. T-R McLaughlin, *A Continuum Limit of the Toda Lattice*, to appear in *Memoirs of the AMS*.
[23] A.S. Fokas, A.R. Its and A.V. Kitaev, Isomonodromic approach in the theory of two-dimensional quantum gravity, *Uspekhi Mat. Nauk* **45** (6) (276) (1990) 135–136 (in Russian).

[24] A.S. FOKAS, A.R. ITS, A.V. KITAEV, Discrete Painlevé equations and their appearance in quantum gravity, *Commun. Math. Phys.* **142** (1991) 313-344.

[25] A.S. Fokas, A.R. Its and A.V. Kitaev, The isomonodromy approach to matrix models in 2D quantum gravity, *Commun. Math. Phys.* **147** (1992) 395–430.

[26] A.S. Fokas, A.R. Its and Xin Zhou, Continuous and discrete Painlevé equations, pp. 33-47 *in* [50].

[27] G. FREUD, *Orthogonale Polynome*, Birkhäuser, 1969 = *Orthogonal Polynomials*, Akadémiai Kiadó/Pergamon Press, Budapest/Oxford, 1971.

[28] G.FREUD, On the coefficients in the recursion formulæ of orthogonal polynomials, *Proc. Royal Irish Acad. Sect. A* **76** (1976), 1-6.

[29] G. Freud, Publications of Géza Freud in approximation theory, *J. Approx. Theory* **46** (1986) 9-15.

[30] J.L. GAMMEL, J. NUTTALL, Note on generalized Jacobi polynomials, *in* "The Riemann Problem, Complete Integrability and Arithmetic Applications" (D. Chudnovsky and G. Chudnovski, Eds.), pp.258-270, Springer-Verlag (Lecture Notes Math. **925**), Berlin, 1982.

[31] W.GAUTSCHI, Computational aspects of orthogonal polynomials, pp.181-216 *in Orthogonal Polynomials: Theory and Practice* (P.NEVAI, editor) *NATO ASI Series C* **294**, Kluwer, Dordrecht, 1990.

[32] W. Gautschi, Algorithm 726: ORTHPOL– A package of routines for generating orthogonal polynomials and Gauss-type quadratures, *ACM Trans. Math. Soft.* **20** (1994) 21-62.

[33] B. Grammaticos, B. Dorizzi, Integrable discrete systems and numerical integrators, *Math. and Computers in Simul.* **37** (1994) 341-352.

[34] B. Grammaticos, A. Ramani, V. Papageorgiou, Do integrable mappings have the Painlevé property? *Phys. Rev. Lett.* **67** (1991) 1825-1828.

[35] F.A. Grünbaum, L. Haine, The $q-$version of a theorem of Bochner, *J. Comp. Appl. Math.* **68** (1996) 103-114.

[36] F.A. Grünbaum, L. Haine, Some functions that generalize the Askey-Wilson polynomials, to appear in *Commun. Math. Phys.*

[37] W.HAHN, On differential equations for orthogonal polynomials, *Funk. Ekvacioj*, **21** (1978) 1-9.

[38] W.HAHN, Über Orthogonalpolynome, die linearen funktionalgleichungen genügen, pp. 16-35 *in Polynômes Orthogonaux et Applications, Proceedings, Bar-le-Duc 1984*, (C.BREZINSKI & al., editors), *Lecture Notes Math.* **1171**, Springer, Berlin 1985.

[39] L. HAINE, E. HOROZOV, Toda orbits of Laguerre polynomials and representations of the Virasoro algebra. *Bull. Sci. Math.* (2) **117** (1993) 485-518.

[40] E. HENDRIKSEN, H. van ROSSUM, A Padé-type approach to non-classical orthogonal polynomials, *J. Math. An. Appl.* **106** (1985) 237-248.

[41] E. HENDRIKSEN, H. van ROSSUM, Semi-classical orthogonal polynomials, pp. 354-361 *in Polynômes Orthogonaux et Applications, Proceedings, Bar-le-Duc 1984*, (C.BREZINSKI & al., editors), *Lecture Notes Math.* **1171**, Springer, Berlin 1985.

[42] J. Hietarinta, Rational solutions to d$-$P$_{\rm IV}$, these Proceedings.

[43] A.R. Its, these Proceedings.

[44] M.KAC, P. van MOERBEKE, On an explicitly soluble system on nonlinear differential equations related to certain Toda lattices, *Adv. Math.* **16** (1975) 160-169.

[45] K. Kajiwara, Y. Ohta, J. Satsuma, B. Grammaticos, A. Ramani, Casorati determinant solutions for the discrete Painlevé-II equation, *J. Phys. A: Math. Gen.* **27** (1994) 915-922.

[46] Y. Kodama, K. T-R McLaughlin, Explicit integration of the full symmetric Toda hierarchy and the sorting property, *Letters in Math. Phys.* **37** (1996) 37-47.

[47] R. Koekoek, R.F. Swarttouw, The Askey-scheme of hypergeometric orthogonal polynomials and its $q-$analogue, Report 94-05, Delft Univ. of Technology, Faculty TWI, 1994. See also the "Askey-Wilson scheme project" entry in http://www.cs.vu.nl/~rene/ for errata and electronic access.

[48] M. Kruskal, Remarks on trying to unify the discrete with the continuous, these Proceedings.

[49] E. LAGUERRE, Sur la réduction en fractions continues d'une fraction qui satisfait à une équation différentielle linéaire du premier ordre dont les coefficients sont rationnels, *J. Math. Pures Appl. (4)* **1** (1885), 135-165 = pp. 685-711 *in Oeuvres*, Vol.II, Chelsea, New-York 1972.

[50] D. LEVI, P. WINTERNITZ, editors: *Painlevé Transcendents. Their Asymptotics and Physical Applications. NATO ASI Series: Series B: Physics* **278**, Plenum Press, N.Y., 1992.

[51] J.S.LEW, D.A.QUARLES, Nonnegative solutions of a nonlinear recurrence, *J. Approx. Th.* **38** (1983), 357-379.

[52] J.-M. LIU, G. MÜLLER, Infinite-temperature dynamics of the equivalent-neighbor XYZ model, *Phys. Rev. A* **42** (1990) 5854-5864.

[53] D.S. LUBINSKY, A survey of general orthogonal polynomials for weights on finite and infinite intervals, *Acta Applicandæ Mathematicæ* **10** (1987) 237-296.

[54] D.S. Lubinsky, An update on general orthogonal polynomials and weighted approximation on the real line, *Acta Applicandæ Math.* **33** (1993) 121–164.

[55] A.P. MAGNUS, Riccati acceleration of Jacobi continued fractions and Laguerre-Hahn orthogonal polynomials, pp. 213-230 *in Padé Approximation and its Applications, Proceedings, Bad Honnef 1983*, Lecture Notes Math. **1071** (H.WERNER & H.T.BÜNGER, editors), Springer-Verlag, Berlin, 1984.

[56] A.P. MAGNUS, A proof of Freud's conjecture about orthogonal polynomials related to $|x|^\rho \exp(-x^{2m})$ for integer m, pp. 362-372 *in Polynômes Orthogonaux et Applications, Proceedings Bar-le-Duc 1984.* (C.BREZINSKI *et al.*, editors), *Lecture Notes Math.* **1171**, Springer-Verlag, Berlin 1985.

[57] A.P. MAGNUS, On Freud's equations for exponential weights, *J. Approx. Th.* **46** (1986) 65-99.

[58] A.P. MAGNUS, Associated Askey-Wilson polynomials as Laguerre-Hahn orthogonal polynomials, pp. 261-278 *in Orthogonal Polynomials and their Applications, Proceedings Segovia 1986.* (M.ALFARO *et al.*, editors), *Lecture Notes Math.* **1329**, Springer-Verlag, Berlin 1988.

[59] A.P. Magnus, Refined asymptotics for Freud's recurrence coefficients, pp. 196-200 *in* V.P. Havin, N.K. Nikolski (Eds.): *Linear and Complex Analysis Book 3, Part II*, Springer Lecture Notes Math. **1574**, Springer, 1994.

[60] A.P.MAGNUS: Painlevé-type differential equations for the recurrence coefficients of semi-classical orthogonal polynomials. *J. Comp. Appl. Math.* **57** (1995) 215-237.

[61] A.P. MAGNUS Asymptotics for the simplest generalized Jacobi polynomials recurrence coefficients from Freud's equations: numerical explorations, *Annals of Numerical Mathematics* **2** (1995) 311-325.

[62] F. MARCELLÁN, I. ALVAREZ ROCHA, On semiclassical linear functionals: integral representations, *J. Comp. Appl. Math.* **57** (1995) 239-249.

[63] F. Marcellán, I.A. Rocha, Complex path integral representation for semiclassical linear functionals, preprint.

[64] P. MARONI, Le calcul des formes linéaires et les polynômes orthogonaux semi-classiques, pp. 279-290 *in Orthogonal Polynomials and their Applications. Proceedings, Segovia 1986* (M. ALFARO *et al.*, editors), *Lecture Notes Math.* **1329**, Springer, Berlin 1988.

[65] P. MARONI, Une théorie algébrique des polynômes orthogonaux. Application aux polynômes orthogonaux semi-classiques. pp. 95-130 *in Orthogonal Polynomials and their Applications* (C.BREZINSKI *et al.*, editors), *IMACS Annals on Computing and Applied Mathematics* **9** (1991), Baltzer AG, Basel.

[66] L.M. Milne-Thomson, *The Calculus of Finite Differences*, Macmillan, London, 1933.

[67] P.NEVAI, Two of my favorite ways of obtaining asymptotics for orthogonal polynomials, pp.417-436 *in Anniversary Volume on Approximation Theory and Functional Analysis*, (P.L.BUTZER, R.L.STENS and B.Sz.-NAGY, editors), **ISNM 65**, Birkhäuser Verlag, Basel, 1984.

[68] P. Nevai, Letter to a friend, *J. Approx. Theory*, **46** (1986) 5-8.

[69] P. NEVAI, Géza Freud, orthogonal polynomials and Christoffel functions. A case study, *J. Approx. Theory* **48** (1986), 3-167.

[70] P.NEVAI, Research problems in orthogonal polynomials, pp. 449-489 *in Approximation Theory VI*, vol. **2** (C.K.CHUI, L.L.SCHUMAKER & J.D.WARD, editors), Academic Press, 1989.

[71] P. Nevai, Introduction to chapter 13 (Orthogonal polynomials) of V.P. Havin, N.K. Nikolski (Eds.), *Linear and Complex Analysis Problem Book 3, Part II, Springer Lecture Notes Math.* **1574**, Springer, 1994.

[72] J.NUTTALL, Asymptotics of diagonal Hermite-Padé polynomials, *J.Approx. Th.* **42** (1984) 299-386.

[73] E.P. O'REILLY, D. WEAIRE, On the asymptotic form of the recursion basis vectors for periodic Hamiltonians, *J. Phys. A: Math. Gen.* **17** (1984) 2389-2397.

[74] V. Papageorgiou, these Proceedings

[75] V. Papageorgiou, B. Grammaticos, A. Ramani, Orthogonal polynomial approach to discrete Lax pairs for initial boundary-value problems of the qd algorithm, *Lett. Math. Phys.* **34** (1995) 91-101.

[76] O.PERRON, *Die Lehre von den Kettenbrüchen*, 2nd edition, Teubner, Leipzig, 1929 = Chelsea,

[77] J.A. SHOHAT, A differential equation for orthogonal polynomials, *Duke Math. J.* **5** (1939),401-417.

[78] W. Van Assche, *Asymptotics for Orthogonal Polynomials, Springer Lecture Notes Math.* **1265**, Springer, 1987.

[79] J.WIMP, Current trends in asymptotics: some problems and some solutions, *J. Comp. Appl. Math.* **35** (1991) 53-79.

Institut de Mathématique Pure et Appliquée, Université Catholique de Louvain, Chemin du Cyclotron,2, B-1348 Louvain-la-Neuve (Belgium)

E-mail address: magnus@anma.ucl.ac.be , www: http://math.ucl.ac.be/~magnus/